



SOCIAL PRESCRIBING

80% non clinical
20% of GP visits non clinical
Green social prescribing treats mild depression
All evaluations show a reduction in health costs
25% cut in general practice consultations
9% cut in Hospital Consultations

SOCIAL PRESCRIBING

Life expectancy stalling

By 2040 1 in 5 living with major illness

Least deprived area 70 years good health

Deprived area only 50 years good health

CHILDREN

National

4.2 million living in poverty (3.6 10 years ago) 62% Bangladeshi Children live below the poverty line

Age 2-10 12% obese

Age 11-15 19% obese

East end on London

50% of 5-year-olds have severe dental caries and vitamin D Deficiency

COVID INEQUALITY

Multigenerational households

Jobs with multiple contact with people

Poorer areas

Low engagement with, and trust of institutions


People who suffer Racism and Discrimination

Antidepressant drug consumption per 1,000 people, 2021




Daily intake of antidepressants per 1,000 people, calculated as the 'defined daily dose'.

This enables us to compare overall levels of drug consumption, accounting for the fact that different antidepressants have different recommended dosages.





Standardised death rate due to intentional self-harm, be (Number of deaths due to intentional self-harm per 100 000 in



Suicide Rates per 100,000 population

and

antidepressants per 1000

	Suicide /Pills
Finland 15.3	81.5
Sweden 14.7	108.9
Norway11.8	61.1





























































